



Contents lists available at ScienceDirect

# Trends in Food Science & Technology

journal homepage: <http://www.journals.elsevier.com/trends-in-food-science-and-technology>

## Commentary

# Novel strategies for improving dietary exposure assessment: Multiple-data fusion is a more accurate measure than the traditional single-biomarker approach



Mar Garcia-Aloy <sup>a, b, 1</sup>, Montserrat Rabassa <sup>a, b, 1</sup>, Patricia Casas-Agustench <sup>a</sup>,  
Nicole Hidalgo-Liberona <sup>a</sup>, Rafael Llorach <sup>a, b</sup>, Cristina Andres-Lacueva <sup>a, b, \*</sup>

<sup>a</sup> Biomarkers and Nutrimental Laboratory, Department of Nutrition, Food Sciences and Gastronomy, Food Technology Reference Net (XaRTA), Nutrition and Food Safety Research Institute (INSA), Campus Torribera, Faculty of Pharmacy and Food Sciences, University of Barcelona, Barcelona, Spain

<sup>b</sup> CIBER de Fragilidad y Envejecimiento Saludable (CIBERFES), Instituto de Salud Carlos III, Barcelona, Spain

## ARTICLE INFO

### Article history:

Received 7 November 2016

Received in revised form

2 March 2017

Accepted 28 April 2017

Available online 29 April 2017

### Keywords:

Dietary assessment

Dietary questionnaires

Biomarkers

Metabolomics

Multi-metabolite biomarker model

Nutrimentalomics

## ABSTRACT

**Background:** Accurate measurement of food intake is the cornerstone of understanding the links between diet and optimal health status or risk of disease. The utilization of metabolomics approaches is revolutionizing the field of dietary assessment by associating metabolic profiles with intake of specific foods or dietary patterns and/or investigating human health status in nutritional trials. Combining dietary biomarkers with conventional dietary assessment methods is considered a potential strategy for tackling the complexity of dietary exposure fingerprinting.

**Scope and approach:** We discuss existing approaches among dietary assessment methods and dietary biomarkers. A combined approach taking into consideration data from dietary questionnaires with measurements of dietary biomarkers is emphasized.

**Key findings and conclusions:** Trends in novel strategies for improving dietary exposure assessment will be influenced by the discovery and validation of dietary exposure biomarkers. Among different strategies, multi-metabolite biomarker panels enable more reliable estimation of dietary exposure than does the traditional single-biomarker approach. Therefore, a combined approach using data from dietary questionnaires along with measurements of dietary biomarkers is considered an excellent strategy for improving dietary exposure assessment.

© 2017 Elsevier Ltd. All rights reserved.

## 1. Introduction

Diet and nutrition are major determinants of human health. Accurate measurement of food intake is the cornerstone of understanding the links between diet and optimal health status or risk of disease. Dietary assessment has been traditionally performed using conventional methodologies of food surveys such as food frequency questionnaires (FFQs), 24-h dietary recalls or food records. However, the accuracy of the dietary intake and nutritional status is frequently challenged due to the subjective nature of these dietary instruments. This limitation can be improved by the

application of metabolomics to characterize dietary exposure.

Metabolomics approaches are revolutionizing the field of dietary assessment by associating metabolic profiles with intake of specific foods or dietary patterns, and/or investigating human health status in nutritional trials. Recently, exploring the food metabolome has been defined as a data-driven strategy for identifying novel biomarkers and improving the accuracy of measurement of dietary exposures by traditional dietary assessment instruments (Scalbert et al., 2014). In this way, the use of metabolomics has enabled the identification of novel and robust biomarkers of food or nutrient intake, which provide an objective measure of exposure that is devoid of many of the biases and errors associated with self-reported methods (Scalbert et al., 2014).

However, there are some factors not present in the traditional dietary assessment instruments that could misrepresent biomarker measures of dietary intake. These factors include genetic variability (e.g. biological variation in nutrient absorption and metabolism,

\* Corresponding author. Biomarkers and Nutrimental Laboratory, Department of Nutrition, Food Sciences and Gastronomy, Faculty of Pharmacy and Food Sciences, University of Barcelona, Av. Joan XXIII s/n, 08028, Barcelona, Spain.

E-mail address: [candres@ub.edu](mailto:candres@ub.edu) (C. Andres-Lacueva).

<sup>1</sup> Equal contribution.

epigenetic variation or gene-gene interactions), lifestyle/physiological factors (e.g. smoking, alcohol consumption, physical exercise or influence of microbiota), dietary factors (e.g. nutrient bioavailability or nutrient-nutrient interactions), biological samples and analytical methodology (Scalbert et al., 2014). Further research will address this issue and identify the best emerging dietary biomarkers. Therefore, as biomarkers cannot replace conventional dietary assessment methods, the use of conventional dietary instruments together with dietary biomarkers is considered the best strategy for tackling the complexity of dietary exposure fingerprinting. Fig. 1 represents a schematic diagram for combining dietary questionnaires with biomarkers.

The main aim of this paper is to highlight existing approaches aimed at improving dietary exposure assessment and the novel combined measure considering data from dietary questionnaires with measurements of dietary biomarkers.

## 2. Dietary assessment methods

### 2.1. Dietary questionnaires

In nutritional studies, the traditional method of collecting data on foods and beverages consumed over a prescribed period of time relies on information gathered using dietary assessment questionnaires. As previously mentioned, the most commonly used dietary assessment methods are open-ended questionnaires such as food records or 24-h dietary recalls, or closed-ended questionnaires including FFQs (Tucker, 2007). Generally, these instruments require a systematic estimation of the frequency and usual serving size of foods, as well as detailed information about the ingredients of a meal recipe, combinations of foods consumed together, and sometimes cooking processes, which may influence the estimation of exposure to a particular dietary constituent. Moreover, the estimation of nutrient/compound intakes depends largely on the existence of appropriate, complete, reliable and up-to-date food composition tables or databases. Many users are often not conscious of the high composition variability, particularly in terms of micronutrients (e.g. vitamins) and phytochemicals, between similar foods, and even in the same type of food (e.g. raw, cooked or processed foods). For this reason, a great dependency on professionals' knowledge is needed to generate, compile, update and use the food composition data adequately. Indeed, the Food and Agriculture Organization of the United Nations (FAO) and the International Network of Food Data Systems (INFOODS) have recently published three comprehensive guidelines on conversions, data evaluation and food matching in order to improve and harmonize the compilation of food composition data, which could also lead to more accurate nutrient intake estimations (Charrondiere et al., 2016).

The consumption data obtained through these methods are used to compute the intake of whole diets/dietary patterns, food groups, foods, energy, nutrients, bioactive compounds and other food components. Recently, dietary pattern analysis has emerged as a useful approach for investigating diet-disease associations. Thus, eating patterns may be more predictive of disease risk than isolated analysis on foods or nutrients (Hu, 2002). The 2015 Dietary Guidelines Advisory Committee recognized the advantages that dietary patterns offer as an approach for informing public health recommendations (U.S. Department of Health and Human Services & U.S. Department of Agriculture, 2015, p. 2). Different approaches for developing dietary patterns exist (Lassale et al., 2016). The most prominent methods are, a priori, numerical indexes, scores that measure adherence to disease-specific dietary and lifestyle guidelines (e.g. Dietary Approaches to Stop Hypertension (DASH)), scores that measure adherence to a regional diet (e.g. Mediterranean Diet

Score) and scores based on nutritional guidelines (e.g. Diet Quality Index International (DQI-I)). Other approaches have been proposed to derive patterns by using all food groups available, such as principal component analysis (PCA), reduced rank regression, partial least-squares regression (PLS), confirmatory factor analysis (CFA) and treelet transform (Hu, 2002; Imamura & Jacques, 2011; Varraso et al., 2012).

The strengths of these methods are the lower relative cost, the ease with which the questionnaires can be completed with the help of a trained interviewer (dietician) or by participants themselves (self-reported questionnaires), and the chance to gather a large amount of dietary data. However, the use of questionnaires is also subject to some limitations (Tucker, 2007). The main one is that they are mostly self-reported, wherein the estimation of food portion size is an important source of errors (perception, conceptualization and memory), which could be inappropriate for some populations (children, obese people, and elderly people with cognitive impairment, among others). Such systematic errors inherent in self-reported data plus random errors (e.g. the accuracy of the food composition tables) can bias the estimation of dietary intake. However, the FFQ is the most common dietary assessment method used to estimate habitual dietary intake (e.g. the previous 12 months) of specific nutrients, dietary exposures related to a certain disease or various dietary components in large-population studies, due to its self- or interviewer-administered and economical machine-readable features. In this context, the use of a previously validated FFQ is an essential requirement for improving the measurement of errors previously mentioned. FFQs should be developed specifically for the research objective because diet may be influenced by participant's characteristics such as ethnicity, culture, dietary habits and lifestyle, among others. Nevertheless, more precise measurements are obtained when using multiple 24-h dietary recalls or dietary records, but only short-term intake (actual intake information over the previous 24 hours) is estimated. In this context, long-term intake can also be estimated if repeated during the year. Recently, it has been suggested that a combination of two different dietary assessment instruments, such as four to six 24-h dietary recalls with a FFQ, could improve estimates of dietary intakes with regard to the methods separately. In this study, the association between diet and disease was statistically significant with food records but not with a FFQ (Carroll et al., 2012). Therefore, if new nutritional studies are designed to include FFQs plus repeat 24-h evaluations, further improvements to minimize their measurement errors might be seen by combining data from the two methods. Our group has recently observed that the highest tertile of total dietary polyphenols, which were estimated using a validated FFQ and an ad hoc database on polyphenol content in foods, was not associated with the risk of cognitive and physical decline, frailty and total mortality, in comparison with the lowest tertile (Rabassa, Cherubini, et al., 2015; Rabassa, Zamora-Ros, et al., 2015, 2016; Zamora-Ros et al., 2013). However, an association with total urinary polyphenols was observed. Moreover these studies have demonstrated the importance of assessing dietary polyphenol exposure whenever possible, using dietary biomarkers and not only through dietary questionnaires.

Despite these strengths, more refined and improved techniques of dietary assessment intake are essential to reduce the limitations of traditional dietary questionnaires and also to reduce the cost associated with the collection and processing of dietary data (Illner et al., 2012; Stumbo, 2013). This is being met with intense methodological research and innovative technologies. Many applications of information and communication technologies are currently under development and validation, and great strides have been made. An example of interactive computer-based techniques is a menu-driven standardized 24-h dietary recall program (called EPIC-




Fig. 1. Schematic diagram for combining dietary questionnaires with biomarkers.

SOFT) developed by the European Prospective Investigation into Cancer and Nutrition study (Slimani et al., 2011). The National Cancer Institute in the US has also developed an interactive computer-based approach but with an Internet-based technology, called the Automated Self-Administered 24-h Dietary Recall, which is based on the Automated Multiple-Pass Method approach (Schatzkin et al., 2009). In addition, mobile phone applications have been released such as Nutricam, which allows users to capture images of foods and verbally describe their items before intake, and then to upload both the image and voice file onto a website for analysis (Rollo, Ash, Lyons-Wall, & Russell, 2011). Another example is a wearable electronic device that resembles a necklace and includes a microphone, camera and other sensors. In this case, dietary intakes are collected from the video recording and are calculated automatically (Sun et al., 2010). However, these methods have not yet been widely implemented in large-population studies due to their related technical issues (data transfer, storage, battery life and others) and methodological difficulties such as self-reporting and higher costs. In addition, certain users are not familiar with innovative technologies or new devices. Despite these limitations, automated versions promise to overcome the labour-intensive and costly coding of the 24-h dietary recalls (Shim, Oh, & Kim, 2014).

## 2.2. Dietary biomarkers

Thus far, individual biomarkers of dietary intake have been used to assess exposure to specific foods or food groups. However, this strategy has important limitations and only in some cases it has been successful. The more relevant limitations can be grouped into a wide distribution of food components (low specificity of biomarkers), high interindividual variation and the microbiota metabolism, among others.

A common approach when the research community looks for a biomarker of intake is first to study the food composition, then try

to identify the possible modifications caused by the host metabolism and finally look for metabolites (e.g. biomarkers) in the biofluid (mostly blood and/or urine). Usually, the food composition is very complex (from the quantitative and qualitative point of view) and many of the compounds are widely distributed in different foods. For instance, most polyphenols are present in a wide range of plant foods, such as chlorogenic acids (e.g. coffee and apple) and flavan-3-ols (e.g. cocoa and tea) (Clifford, 2000; Monagas et al., 2010). Another example of a compound widely distributed in many plant foods is vitamin C. This compound has been used as a biomarker of consumption of fruits and vegetables, although differences in concentration between different foods reduces their ability to be a good biomarker (Scalbert et al., 2014). Therefore using the single-biomarker strategy compromises its usefulness because there are a number of factors that limit the prediction of dietary exposure.

It is worth noting that in some cases, similar compounds from different food sources could produce the same biomarker. One of the most relevant cases is ellagitannins. This class of polyphenols (with some differences) is present in foods such as pomegranate, strawberries and walnuts (Espín, Larrosa, García-Conesa, & Tomás-Barberán, 2013). They are poorly absorbed and when they reach the gut they are largely metabolized by the microbiota, producing urolithin derivatives (Espín et al., 2013). Selma, Beltrán, García-Villalba, Espín, & Tomás-Barberán (2014) showed the ability of the bacteria *Gordinobacter* to produce urolithins from ellagitannins. This means that urolithins are biomarkers of ellagitannin intake instead of particular foods. This relevance of the microbial effect is crucial because a significant number of food components are degraded by the colonic microbiota and after absorption and distribution are excreted in urine. Therefore, the real biomarker of intake is provided by the microbiota instead of the host metabolism.

Procyanidins are a good example of this behaviour. This class of

polyphenols is present in many dietary sources, such as cocoa, tea, wine and apples (Monagas et al., 2010). These polyphenols show low bioavailability. However, gut microbiota have the capacity to degrade these metabolites and produce other compounds called hydroxyphenylvalerolactones and hydroxyphenylvaleric acids (Monagas et al., 2010). In fact, these metabolites have been used as biomarkers of procyanidin-rich foods. Both urolithins and hydroxyphenylvalerolactones have been used as biomarkers of intake of single foods (e.g. walnuts and tea, respectively). However, taking into account the variety of dietary sources that could provide these parent compounds (e.g. ellagitannins and procyanidins), these compounds are not suitable for use as single accurate biomarkers.

In relation to the interindividual variation, there are some interesting examples. According to Lars O. Dragsted (2010), creatinine can be considered a potential marker of meat intake (even cooked). However, endogenous levels connected with creatine turnover showed important variations between subjects (Dragsted, 2010). Another example is a recent study about polyphenols and their urinary quantification (Achaintre et al., 2016). In this paper the authors showed that in urine from 475 EPIC participants a total of 34 polyphenols were evaluated and in general these compounds showed large interindividual variations (Achaintre et al., 2016).

There are several examples where a single metabolite could be a potential biomarker of a particular food intake. Some metabolites have been proposed as a biomarker of intake of a particular group of plant foods. However, in these groups there are particular foods that could represent a very important part of the dietary source (e.g. citrus fruits and oranges, or cruciferous vegetables and broccoli). The compound termed proline betaine has been identified in both intervention and cohort studies (Pujos-Guillot et al., 2013) as a candidate biomarker of citrus intake and, in particular, a powerful biomarker of orange intake (Lloyd, Beckmann, Favé, Mathers, & Draper, 2011). This biomarker was detected and its urinary excretion kinetics reported after an intervention study with orange juice consumption (Heinzmann et al., 2010). However, increased urinary excretions of proline betaine have also been observed after diets enriched with rye bran, bringing its specificity under the spotlight (Pekkinen et al., 2015). With regard to cruciferous intake, and in particular that of broccoli, sulforaphane (mainly its mercapturic acid derivative) was proposed as a potential biomarker of intake of this particular food (Dominguez-Perles et al., 2014; Vermeulen, van Rooijen, & Vaes, 2003).

### 2.2.1. Multi-metabolite biomarker models

Given the already mentioned inconsistencies that can limit the usefulness of single biomarkers for dietary intake evaluation, the question emerges of whether a combination of food-derived metabolites, namely a multi-metabolite biomarker panel (MBP), would be more likely to capture dietary exposure and improve the accuracy and precision of dietary assessment. The rationale behind the use of MBPs is that a wider range of metabolites would improve the measurement of dietary intake, capturing a broader perspective of the diet and thereby giving a more complete coverage of dietary exposure. It opens a new framework in the research area of nutritional biomarkers. However, while almost all studies investigating dietary biomarkers have focused on single candidate biomarkers, MBPs have remained practically unexplored and only a few research groups have addressed this question during the last few years. These studies are summarized in Tables 1 and 2.

Over the last few years, our research group has made great efforts to instigate a novel approach for improving dietary exposure assessment through MBPs. This concept has been used in a number of recent studies where we suggested MBPs of walnuts (Garcia-Aloy et al., 2014), wine (Vázquez-Fresno et al., 2015), cocoa (Garcia-Aloy, Llorach, Urpi-Sarda, Jáuregui, et al., 2015) and bread (Garcia-Aloy,

Llorach, Urpi-Sarda, Tulipani, et al., 2015) using urine samples analysed by an untargeted metabolomics approach, and dietary data from FFQs, in studies with different designs. The results showed that MBPs perform better in terms of predicting dietary exposure. Our first study in this field identified an MBP that was highly predictive of walnut intake with an area under the curve [AUC (95% CI)] of 93.4% (90.1–96.8%) and 90.2% (85.9–94.6%) in the training and validation sets, respectively (Garcia-Aloy et al., 2014). In line with these results, a “tartrate-ethyl glucuronide” model showed an AUC of 90.7% (84.5–96.4%) in the training set composed of samples from volunteers that participated in a controlled clinical trial with a nutritional intervention with wine, and an AUC of 92.4% (84.1–100%) in the validation set composed of samples assessed at baseline from a subcohort of volunteers included in the PREDIMED study with a reported wine intake of  $\geq 180$  mL/day (Vázquez-Fresno et al., 2015). Additionally, this model showed promising performance in terms of its sensitivity, which enabled discernment of an intake of one glass of wine 3 days after consumption in an observational study. Another MBP was highly predictive for cocoa consumption [AUC = 95.7% (89.8–100%) in the training set, and 92.6% (81.9–100%) in the validation set]. It was built with one component of the theobromine metabolism (7-methylxanthine) together with another from microbial metabolism of polyphenols (5-(3',4'-dihydroxyphenyl)-valerolactone glucuronide). Both metabolites have been proposed as biomarkers of cocoa intake in studies with different designs (i.e., acute interventions, long-term intervention trials and observational studies) and provided the model with complementary information about habitual cocoa intake (Garcia-Aloy et al., 2015). Finally, an additional MBP was highly predictive for wholegrain bread intake [AUC = 93.1% (88.7–97.4%) and 93.7% (89.4–98.1%) for data from positive and negative ionization mode, respectively], while the MBP designed to evaluate white bread consumption had a reasonably good predictive ability [AUC = 80.6% (72.1–89.0%) and 77.8% (69.1–86.4%) for data from positive and negative ionization mode, respectively] (Garcia-Aloy et al., 2015).

Previously, Campbell et al. (1994) published one of the first studies suggesting a combination of biomarkers of intake. They used a stepwise logistic regression analysis to assess fruit and vegetable consumption. The resultant prediction model included compounds measured in biological samples (three carotenoids determined in plasma) and data from dietary questionnaires (energy intake) (Gross et al., 1994). Later, Nielsen, Freese, Kleemola, and Mutanen (2002) proposed measuring the sum of different flavonoids determined in urine for examining the intake of fruits and vegetables (Nielsen et al., 2002), an approach also suggested in other studies (Brevik, Rasmussen, Drevon, & Andersen, 2004; Krogholm, Haraldsdóttir, Knuthsen, & Rasmussen, 2004). In the same vein, summing individual resveratrol or alkylresorcinol metabolites has also been attempted for assessing wine (Queipo-Ortuño et al., 2012; Rotches-Ribalta et al., 2012; Zamora-Ros et al., 2009) and wholegrain wheat and rye consumptions (Kristensen et al., 2012; Landberg et al., 2009; Linko-Parvinen, Landberg, Tikkanen, Adlercreutz, & Peñalvo, 2007), respectively. In parallel, the ratio between to alkylresorcinols, C17:0/C21:0, has shown the ability to discern between wholegrain wheat and rye intakes (Landberg et al., 2009; Linko-Parvinen et al., 2007). However, although these later examples exhibited a good predictive capacity, these statistical approaches could not give the real weight of each metabolite within the biomarker panel, and therefore more sophisticated approaches could be required for the assessment of dietary exposures for more complex foods, food groups or dietary patterns. At the same time, it is important to bear in mind that using metabolites from the same class could not deal with the problem of specificity previously highlighted.

As mentioned above, recent work from our laboratory applied a

**Table 1**  
Summary of multi-metabolite biomarker panels identified using untargeted metabolomics approaches.

Food item	Statistical test	Metabolites in the panel	Study design	TS & VS Panel-diet associations	Was the panel better than single biomarkers?	Reference
Walnuts	Stepwise logistic regression	3-Indolecarboxylic acid glucuronide 10-Hydroxy-decane-4,6-dienoic acid sulphate Urolithin A glucuronide Tridecadienoic/tridecynoic acid glucuronide Urolithin A sulphate	Observational (cross-sectional)	Yes AUC (95% CI): • TS = 93.4% (90.1–96.8%) • VS = 90.2% (85.9–94.6%)	Yes	<a href="#">Garcia-Aloy et al., 2014</a>
Cocoa	Stepwise logistic regression	7-Methylxanthine 5-(3',4'-dihydroxyphenyl)-valerolactone GlcA	Observational (cross-sectional)	Yes AUC (95% CI): • TS = 95.7% (89.8–100%) • VS = 92.6% (81.9–100%)	Yes	<a href="#">Garcia-Aloy, Llorach, Urpi-Sarda, Jáuregui, et al., 2015</a>
Bread	Stepwise logistic regression	<u>White Bread:</u> HPAA GlcA HMBOA Riboflavin <u>Wholegrain Bread:</u> HHPAA HPPA HMBOA Enterolactone GlcA Pyrraline 3-Indolecarboxylic acid GlcA Riboflavin	Observational (cross-sectional)	No AUC (95% CI): from 77.8% (69.1–86.4%) to 93.7% (89.4–98.1%).	Yes	<a href="#">Garcia-Aloy, Llorach, Urpi-Sarda, Tulipani, et al., 2015</a>
Wine	Stepwise logistic regression	Tartrate Ethyl glucuronide	Sustained intervention [TS] Observational (cross-sectional) [VS]	Yes AUC (95% CI): • TS = 90.7% (84.5–96.4%) • VS = 92.4% (84.1–100%)	Yes	<a href="#">Vázquez-Fresno et al., 2015</a>
Orange juice	Random forest	Stachydrine Methyl glucopyranoside ( $\alpha+\beta$ ) Dihydroferulic acid Galactonate	Sustained intervention	Yes AUC (95% CI): 99.6% (96–100%) Accuracy: • Entire data set = 93% • Hold-out data set = 87.5% • Permutation test: Entire data set: p-value = 0.006 • Hold-out dataset: p-value = 0.004	Yes for some metabolites	<a href="#">Rangel-Huerta et al., 2017</a>
Coffee	Support vector machines	Cyclo(isoleucylprolyl) 1-Methylxanthine Trigonelline	Observational (cross-sectional)	Yes AUC (95% CI): 98% (93–100%)	Yes	<a href="#">Rothwell et al., 2014</a>
Sugar-sweetened beverages	PLS regression	Formate Citrulline Taurine Isocitrate	Observational [TS] Acute intervention [VS]	Yes AUC [TS]: 80% Specificity [TS]: 80% Sensitivity [TS]: 70%	Yes	<a href="#">Gibbons et al., 2015</a>
Nordic diet	PLS-DA	(2-Oxo-2,3-dihydro-1H-indol-3-yl)acetic acid 6-Amino-5-[N-methylformylamino]-1-methyluracil Hydroquinone GlcA 3,4,5,6-Tetrahydrohippurate 3-Indoleacetic acid GlcA Limonene-1,2-diol GlcA Limonene-8,9-diol-GlcA p-Menth-1-2ne-6,8,9-triol Trimethylamine N-oxide Cyclo(Pro-Val) Hippuric acid Octanoyl GlcA Proline betaine 7-Methyluric acid 3,7-Dimethyluric acid 7-Methylxanthine	Sustained intervention	Yes <sup>a</sup> Misclassified samples: • TS: 35% • VS: 19%	NR	<a href="#">Andersen et al., 2014</a>

Diet in general	PCA	Observational (cross-sectional)	No	Correlations	NR	O'Gorman et al., 2014
Perillic acid G1cA						
Perillic acid-8,9-diol-G1cA						
Dihydroperillic acid G1cA						
Pyroglutamyl proline						
Pyraline						
Theobromine						
Unknowns						
LP1:		LP6:		• LP1:		
PCaeC36:2		PEaaC36:1		– SFA: p-value = 0.015		
PCaeC38:3		PSaaC36		– MUFA: p-value = 0.011		
PEaaC22:2		PEaaC38:4:2		– PUFA: p-value = 0.018		
PEaaC34:0				• LP6:		
PEaaC40:4				– Meat: p-value = 0.024		
SMC15:0				– Fish: p-value = 0.026		
SMC19:0				– Vegetable intake: p = 0.036		
SMC20:2				AUCs:		
SMC21:0				• LP1: Fat = 82%		
SMC21:1				• LP6:		
LPcaC18:2				– Meat and vegetable <70%		
LPecC18:0				– Fish = 76%		

AUC, area under the curve; CI, confidence interval; DA, discriminant analysis; G1cA, glucuronide; HHPAA, 2-hydroxy-N-(2-hydroxyphenyl) acetamide; HMBOA, 2-hydroxy-7-methoxy-2H-1,4-benzoxazin-3-one; HPAA, N-(2-hydroxyphenyl) acetamide; HPPA, 2-hydroxy-N-(2-hydroxyphenyl) acetamide; LP, lipid pattern; NR, not reported; PCA, principal component analysis; PLS, partial-least squares; TS, training set; VS, validation set.  
<sup>a</sup> Some samples of the training and validation sets were from the same subjects.

multivariate statistical approach to link dietary data with both targeted and untargeted metabolomics data to identify a series of MBPs (Garcia-Aloy et al., 2014, 2015, 2015; Urpi-Sarda et al., 2015; Vázquez-Fresno et al., 2015). In this approach, stepwise logistic regression analysis was used to include more than one metabolite in biomarker panels and regressed against dietary data to identify MBPs. Other research groups also applied regression analysis for addressing this issue (Jin et al., 2014; McGrath et al., 2016; Neuhaus et al., 2003), whereas, more recently, other multivariate statistical strategies such as PCA, PLS, random forest and support vector machine algorithms have also been used to build MBPs for assessing the consumption of different foods or dietary patterns (Andersen et al., 2014; Gibbons et al., 2015; Marklund et al., 2014; O'Gorman et al., 2014; Rangel-Huerta et al., 2017; Rothwell et al., 2014). These multivariate methods may be of interest in terms of the further discovery of dietary exposure biomarkers. For example, Marklund et al. (2014) created a novel dietary biomarker score based on a combination of several individual biomarker concentrations, using rank scores or principal component analysis to assess compliance in a dietary randomized controlled trial, Systems Biology in Controlled Dietary Interventions and Cohort Studies (SYSDIET), and to investigate how a healthy Nordic diet influences cardiometabolic risk factors (Marklund et al., 2014). Their results suggest that this dietary biomarker score provides a better reflection of the dietary intake and thereby increases the strength of detecting potential cardiometabolic health effects. In addition, they proposed that future studies should evaluate the combined use of dietary biomarkers and reported dietary intake methods for assessing compliance.

Nevertheless, since until now MBPs have been proposed for a limited number of foods, food groups and dietary patterns, further work in this line is required. This will enable a broad coverage of dietary exposure, including the assessment of exposure to bioactive compounds, foods, food groups or complex dietary patterns reflective of habitual dietary intake. The challenge that lies ahead will be finding the simplest combination of metabolites that is able to properly evaluate dietary exposure. In the same vein, the capacity of metabolomics to measure simultaneously a high number of metabolites offers a great opportunity to propose new MBPs clustering in the same model metabolites of different classes, providing the panels with complementary information about dietary intake.

However, before translating MBPs into nutritional epidemiology, appropriate validation steps in the post-discovery phase are essential in order to assess their robustness. Firstly, it is necessary to develop analytical methods for accurate and specific quantification of the metabolites included in each MBP, as recently suggested for simultaneous quantification of multiple biomarkers associated with alcohol intake (Monošík & Dragsted, 2016). Additionally, the specificity, sensibility, kinetics and dose-response relationships of proposed MBPs should be investigated in separate studies with different designs and populations, including people with different genetic and dietary backgrounds.

### 2.3. Methods of combining dietary questionnaires and biomarkers

Taking into account all of the above, and with a view to evaluating the diet-disease relationship, the use of validated dietary biomarkers helps to improve precision in the measure of dietary intake of different foods, nutrients and other components, and to then strengthen the ability to detect dietary effects. In addition, a combined approach using data from dietary questionnaires with measurements of dietary biomarkers has emerged as a good strategy, especially when the extent to which the biomarker mediates the dietary effect is unknown. Freedman, Kipnis, et al.

**Table 2**  
Summary of multi-metabolite biomarker panels identified using targeted approaches.

Food item	Statistical test	Metabolites in the panel	Study design	TS & VS Panel-diet associations	Was the panel better than single biomarkers?	Reference
Wholegrain wheat and rye	Sum Ratio	Sum of AR homologues (AR C17:0-C25:0) AR C17:0/C21:0 ratio	Sustained intervention	– Increase: p-value < 0.05	NR	<a href="#">Linko-Parvinen et al., 2007</a>
Rye wholegrain/bran	Sum Ratio	Sum of AR homologues (C17:0, C19:0, C21:0, C23:0, C25:0) AR C17:0/C21:0 ratio	Sustained intervention	– Increase: p-value < 0.0001	NR	<a href="#">Landberg et al., 2009</a>
Wholegrain wheat	Sum	Sum of AR homologues (C17:0, C19:0, C21:0, C23:0, C25:0)	Sustained intervention	– Increase: p-value < 0.001	NR	<a href="#">Kristensen M, Toubro S, Jensen MG, Ross AB, Riboldi G, Petronio M, Bügel S, &amp; Tetens I, 2012</a>
Wine	Sum	<i>cis</i> -Resveratrol-3- <i>O</i> -GlcA <i>trans</i> -Resveratrol-3- <i>O</i> -GlcA <i>cis</i> -Resveratrol-4'- <i>O</i> -GlcA <i>trans</i> -Resveratrol-3- <i>O</i> -sulphate <i>cis</i> -Resveratrol-4'- <i>O</i> -sulphate <i>trans</i> -Resveratrol-4'- <i>O</i> -sulphate	Observational (cross-sectional)	– Correlation: $r = 0.895$ (p-value < 0.001) AUC (95% CI) = 98.3% (97.3–99.0%) Sensitivity (95% CI) = 93.3% (91.5–94.7%) Specificity (95% CI) = 92.1% (90.2–93.7%)	NR	<a href="#">Zamora-Ros et al., 2009</a>
	Sum	<u>Resveratrol Metabolites:</u> <i>cis</i> -Resveratrol-3- <i>O</i> -GlcA <i>trans</i> -Resveratrol-3- <i>O</i> -GlcA <i>cis</i> -Resveratrol-4'- <i>O</i> -GlcA <i>cis</i> -Resveratrol-3- <i>O</i> -sulphate <i>trans</i> -Resveratrol-3- <i>O</i> -sulphate <i>cis</i> -Resveratrol-4'- <i>O</i> -sulphate <i>trans</i> -Resveratrol-4'- <i>O</i> -sulphate	Sustained intervention	– Increase: p-value < 0.05	NR	<a href="#">Queipo-Ortuño et al., 2012</a>
	Sum	<u>Dihydroresveratrol Metabolites</u> <u>Total Metabolites (Resveratrol and Dihydroresveratrol)</u> <u>Resveratrol Phase II Metabolites</u> <i>cis</i> -Resveratrol-3- <i>O</i> -GlcA <i>trans</i> -Resveratrol-3- <i>O</i> -GlcA <i>cis</i> -Resveratrol-4'- <i>O</i> -GlcA <i>trans</i> -Resveratrol-4'- <i>O</i> -GlcA <i>cis</i> -Resveratrol-3- <i>O</i> -sulphate <i>trans</i> -Resveratrol-3- <i>O</i> -sulphate <i>cis</i> -Resveratrol-4'- <i>O</i> -sulphate <i>trans</i> -Resveratrol-4'- <i>O</i> -sulphate <i>trans</i> -Resveratrol-3,4'- <i>O</i> -disulphate Resveratrol sulphoglucuronide <u>Resveratrol Glucosides</u> <i>cis</i> -Piceid <i>trans</i> -Piceid Piceid-GlcA Piceid sulphate Gut microbial metabolism of Resveratrol Dihydroresveratrol Dihydroresveratrol-GlcA Dihydroresveratrol-sulphate Dihydroresveratrol-sulphoglucuronide	Sustained intervention	– Increase: p-value < 0.05	NR	<a href="#">Rotches-Ribalta et al., 2012</a>
		<u>Urine - hydrolysed samples:</u> 2,4-Dihydroxybenzoic acid Gallic acid Ethylgallate	Sustained intervention	Yes AUCs (95% CI):	Yes for most MBP	<a href="#">Uрпи-Sarda et al., 2015</a>

	Stepwise logistic regression	<u>Urine - non-hydrolysed samples:</u> Methylgallic acid sulphate Ethylgallate sulphate <u>Plasma - hydrolysed samples:</u> 3-Hydroxyphenylacetic acid Gallic acid <i>p</i> -Coumaric acid				<ul style="list-style-type: none"> <li>Urine: from 96.00% (89.24–100%) to 98.68% (97.13–100%)</li> <li>Plasma: from 80.13% (71.75–88.51%) to 91.07% (80.22–100%)</li> </ul> Increase: <i>p</i> -value < 0.001	NR	Brevik et al., 2004	
Fruit & vegetables	Sum	Eriodictyol Naringenin Hesperetin Quercetin	Kaempferol Isorhamnetin Tamarixetin Phloretin	Sustained intervention	–				
				Acute intervention	–	Correlation: • 24 h urine: $r = 0.86$ ( $p$ -value < 0.000001) • morning urine: $r = 0.43$ ( $p$ -value < 0.01)	Yes for 24 h urine No for morning urine	Krogholm et al., 2004	
				Sustained intervention	–	Correlation: $r = 0.35$ ( $p$ -value = 0.0007) Increase: $p$ -value < 0.0001	Yes for most single biomarkers	Nielsen et al., 2002	
	Stepwise logistic regression	Vitamin C b-Carotene Lutein		Observational (cross-sectional)	–	Correlation: $r = 0.42$	Yes		Cooper et al., 2015
				Observational (cross-sectional)	No	Explained variability: 53%	NR		Gross et al., 1994
Regression model	$\alpha$ -Carotene Energy intake Lutein $\beta$ -Cryptoxanthin Vitamin C Carotenoids (cholesterol-adjusted) Ferric-reducing antioxidant power		Sustained intervention	Yes <sup>a</sup>	Correlation • TS: $r = 0.47$ ( $p$ -value < 0.001) • VS: $r = 0.18$ –0.36 ( $p$ -value $\leq 0.05$ )	Yes		Jin et al., 2014	
Logistic regression	Vitamin C Carotenoids		Sustained intervention	No	Correct allocation: 45–86%	Yes for most studies		McGrath et al., 2016	
Diet Quality Index Score	Stepwise linear regressions	Vitamin C $\alpha$ -Tocopherol $\alpha$ -Carotene	$\beta$ -Cryptoxanthin Oleic acid Stearic acid	Observational (cross-sectional)	No	NR	NR	Neuhouser, Patterson, King, Horner, & Lampe, 2003	
Mediterranean diet	NR	Carotenes $\beta$ Vitamin E	EPA DHA	Observational (cross-sectional)	No	Correlation: $r = -0.52$ ( $p$ -value = 0.03)	NR	Gerber, 2006	
Nordic diet	Rank scores PCA	$\alpha$ -Linolenic acid $\beta$ -Carotene Alkylresorcinols	EPA DHA	Sustained intervention	No	NR	NR	Marklund et al., 2014	

AR, alkylresorcinol; AUC, area under the curve; CI, confidence interval; DHA, docosahexaenoic acid; EPA, eicosapentaenoic acid; GlcA, glucuronide; MBP, multi-metabolite biomarker panel; NR, not reported; PCA, principal component analysis; TS, training set; VS, validation set.

<sup>a</sup> Samples of the training and validation sets were from the same subjects.



(2010) and Freedman, Tasevska, et al. (2010) proposed two main direct approaches, Howe's method with ranks or PCA, for combining questionnaires and biomarkers related to whole diet, food group, food, nutrient or other food components (Freedman, Kipnis, et al., 2010; Freedman, Tasevska, et al., 2010). Comparing the results obtained from the two approaches, Freedman, Kipnis, et al. (2010) and Freedman, Tasevska, et al. (2010) found that Howe's method gives results close to those from the PCA approach (Freedman, Kipnis, et al., 2010; Freedman, Tasevska, et al., 2010). It can be said that Howe's method is more appropriate due to its simplicity than the approach based on PCA. Later, these authors proposed a more complex modelling-based approach, namely the regression calibration method, for combining dietary biomarkers and reports that recovers lost power and gives unbiased relative risk estimates (Freedman et al., 2011). However, this method is not applicable to food patterns or foods that have no known specific biomarkers. Unlike this, PCA and Howe's method do not require knowledge of the quantitative relationship between biomarker level and true usual intake. Freedman, Kipnis, et al. (2010) and Freedman, Tasevska, et al. (2010) applied Howe's method in an analysis of a diet-disease association in a real example from the Carotenoids and Age-Related Eye Disease Study (CAREDS) (Freedman, Kipnis, et al., 2010; Freedman, Tasevska, et al., 2010). In their study, the estimated odds ratios [OR (95% CI)] for the primary nuclear cataract outcome using the reported dietary intake (FFQ-lutein plus zeaxanthin), the biomarker level (serum lutein plus zeaxanthin) and the combined FFQ-biomarker were 0.77 (0.57–1.02), 0.69 (0.51–0.94) and 0.66 (0.48–0.91), respectively. As can be seen from the statistical significance, the combination of exposure was higher than that for the FFQ and biomarker alone. In addition, an increase in statistical power was also observed in detecting a diet-disease association. Recently, Rabassa, Cherubini, et al. (2015) and Rabassa, Zamora-Ros, et al. (2015) applied this approach to study the association between habitual dietary resveratrol exposure and the development of frailty syndrome in older adults from the INCHIANTI study (Rabassa et al., 2015). Inverse associations between resveratrol exposure and frailty syndrome risk were observed for FFQ-total resveratrol [OR for comparison of extreme tertiles = 0.17 (0.05–0.63)], biomarker-total urinary resveratrol [0.32 (0.09–1.11)] and FFQ&biomarker-total resveratrol [0.11 (0.03–0.45)]. The most successful results from the combined exposure measure will emerge when the strength of the associations for each separate exposure is similar, as occurs in the examples described in this section.

### 3. Future trends

Future research trends should focus on exploring more novel approaches for the discovery and validation of dietary exposure biomarkers. While considerable progress has been made in demonstrating how dietary biomarkers can be used as dietary assessment tools, a number of challenges have still to be overcome before they can achieve their complete validation, including both analytical and biological perspectives. Recent studies have suggested different strategies for identifying panels of biomarkers. They have also demonstrated that MBPs offer a more reliable estimation of dietary exposure than the traditional single-biomarker approach used until now. Therefore, future studies will face up to the complexity of evaluating the use of MBPs in combination with conventional dietary questionnaires to assess dietary exposure fingerprinting. These advances will enable more detailed information to be obtained about the associations between diet and health, providing better evidence of the development of health claims and dietary advice for both public health institutions and the food industry.

### Conflict of interest

The authors have declared no conflict of interest.

### Acknowledgements

This work was supported by the Spanish Ministry of the Economy and Competitiveness (MINECO) together with the *Joint Programming Initiative* "A Healthy Diet for a Healthy Life" (JPI HDHL, website: <http://www.healthydietforhealthylife.eu>) [grant number FOOTBALL-PCIN-2014-133]; funds from CIBERFES (co-funded by the FEDER Program from EU); and through the EIT Health innovation project COOK2HEALTH. EIT Health is supported by the European Institute of Innovation and Technology, a body of the European Union. We thank the award of the Generalitat de Catalunya's Agency AGAUR [grant number 2014SGR1566].

### References

- Achaintre, D., Buleté, A., Cren-Olivé, C., Li, L., Rinaldi, S., & Scalbert, A. (2016). Differential isotope labeling of 38 dietary polyphenols and their quantification in urine by liquid chromatography electrospray ionization tandem mass spectrometry. *Analytical Chemistry*, 88, 2637–2644.
- Andersen, M.-B. S., Rinnan, Å., Manach, C., Poulsen, S. K., Pujos-Guillot, E., Larsen, T. M., et al. (2014). Untargeted metabolomics as a screening tool for estimating compliance to a dietary pattern. *Journal of Proteome Research*, 13, 1405–1418.
- Brevik, A., Rasmussen, S. E., Drevon, C. A., & Andersen, L. F. (2004). Urinary excretion of flavonoids reflects even small changes in the dietary intake of fruits and vegetables. *Cancer Epidemiology and Prevention Biomarkers*, 13.
- Carroll, R. J., Midthune, D., Subar, A. F., Shumakovich, M., Freedman, L. S., Thompson, F. E., et al. (2012). Taking advantage of the strengths of 2 different dietary assessment instruments to improve intake estimates for nutritional epidemiology. *American Journal of Epidemiology*, 175, 340–347.
- Charondiere, U. R., Rittenschober, D., Nowak, V., Stadlmayr, B., Wijesinha-Bettoni, R., & Haytowitz, D. (2016). Improving food composition data quality: Three new FAO/INFOODS guidelines on conversions, data evaluation and food matching. *Food Chemistry*, 193, 75–81.
- Clifford, M. N. (2000). Chlorogenic acids and other cinnamates – nature, occurrence dietary burden, absorption and metabolism. *Journal of the Science of Food and Agriculture*, 80, 1033–1043.
- Cooper, A. J. M., Sharp, S. J., Luben, R. N., Khaw, K.-T., Wareham, N. J., & Forouhi, N. G. (2015). The association between a biomarker score for fruit and vegetable intake and incident type 2 diabetes: the EPIC-Norfolk study. *European Journal of Clinical Nutrition*, 69, 449–454.
- Dominguez-Perles, R., Medina, S., Moreno, D.Á., García-Viguera, C., Ferreres, F., & Gil-Izquierdo, Á. (2014). A new ultra-rapid UHPLC/MS/MS method for assessing glucoraphanin and sulforaphane bioavailability in human urine. *Food Chemistry*, 143, 132–138.
- Dragsted, L. O. (2010). Biomarkers of meat intake and the application of nutrigenomics. *Meat Science*, 84, 301–307.
- Espín, J. C., Larrosa, M., García-Conesa, M. T., & Tomás-Barberán, F. (2013). Biological significance of urolithins, the gut microbial ellagic acid-derived metabolites: The evidence so far. *Evidence-based Complementary and Alternative Medicine*, 2013, 270418.
- Freedman, L. S., Kipnis, V., Schatzkin, A., Tasevska, N., & Potischman, N. (2010). Can we use biomarkers in combination with self-reports to strengthen the analysis of nutritional epidemiologic studies? *Epidemiologic Perspectives & Innovations*, 7, 2.
- Freedman, L. S., Tasevska, N., Kipnis, V., Schatzkin, A., Mares, J., Tinker, L., et al. (2010). Gains in statistical power from using a dietary biomarker in combination with self-reported intake to strengthen the analysis of a diet-disease association: An example from CAREDS. *American Journal of Epidemiology*, 172, 836–842.
- Freedman, L. S., Midthune, D., Carroll, R. J., Tasevska, N., Schatzkin, A., Mares, J., et al. (2011). Using regression calibration equations that combine self-reported intake and biomarker measures to obtain unbiased estimates and more powerful tests of dietary associations. *American Journal of Epidemiology*, 174, 1238–1245.
- García-Aloy, M., Llorach, R., Urpi-Sarda, M., Jáuregui, O., Corella, D., Ruiz-Canela, M., et al. (2015). A metabolomics-driven approach to predict cocoa product consumption by designing a multimetabolite biomarker model in free-living subjects from the PREDIMED study. *Molecular Nutrition & Food Research*, 59, 212–220.
- García-Aloy, M., Llorach, R., Urpi-Sarda, M., Tulipani, S., Estruch, R., Martínez-González, M. A., et al. (2014). Novel multimetabolite prediction of walnut consumption by a urinary biomarker model in a free-living population: The PREDIMED study. *Journal of Proteome Research*, 13, 3476–3483.
- García-Aloy, M., Llorach, R., Urpi-Sarda, M., Tulipani, S., Salas-Salvadó, J., Martínez-

- González, M. A., et al. (2015). Nutrimetabolomics fingerprinting to identify biomarkers of bread exposure in a free-living population from the PREDIMED study cohort. *Metabolomics*, 11, 155–165.
- Gerber, M. (2006). Qualitative methods to evaluate Mediterranean diet in adults. *Public Health Nutrition*, 9, 147–151.
- Gibbons, H., McNulty, B. A., Nugent, A. P., Walton, J., Flynn, A., Gibney, M. J., et al. (2015). A metabolomics approach to the identification of biomarkers of sugar-sweetened beverage intake. *The American Journal of Clinical Nutrition*, 101, 471–477.
- Gross, M. D., Potter, J. D., Campbell, D. R., Martini, M. C., Slavina, J. L., & Grandits, G. A. (1994). Plasma carotenoids as biomarkers of vegetable and fruit intake. *Cancer Epidemiology Biomarkers and Prevention*, 3, 493–500.
- Heinzmann, S. S., Brown, I. J., Chan, Q., Bictash, M., Dumas, M. E., Kochhar, S., et al. (2010). Metabolic profiling strategy for discovery of nutritional biomarkers: Proline betaine as a marker of citrus consumption. *The American Journal of Clinical Nutrition*, 92, 436–443.
- Hu, F. B. (2002). Dietary pattern analysis: A new direction in nutritional epidemiology. *Current Opinion in Lipidology*, 13, 3–9.
- Illner, A.-K., Freisling, H., Boeing, H., Huybrechts, I., Crispim, S. P., & Slimani, N. (2012). Review and evaluation of innovative technologies for measuring diet in nutritional epidemiology. *International Journal of Epidemiology*, 41, 1187–1203.
- Imamura, F., & Jacques, P. F. (2011). Invited commentary: Dietary pattern analysis. *American Journal of Epidemiology*, 173, 1105–8–10.
- Jin, Y., Gordon, M. H., Alimbetov, D., Chong, M. F.-F., George, T. W., Spencer, J. P., et al. (2014). A novel combined biomarker including plasma carotenoids, vitamin C, and ferric reducing antioxidant power is more strongly associated with fruit and vegetable intake than the individual components. *The Journal of Nutrition*, 144, 1866–1872.
- Kristensen, M., Toubro, S., Jensen, M. G., Ross, A. B., Riboldi, G., Petronio, M., et al. (2012). Whole grain compared with refined wheat decreases the percentage of body fat following a 12-week, energy-restricted dietary intervention in postmenopausal women. *The Journal of Nutrition*, 142, 710–716.
- Krogholm, K. S., Haraldsdóttir, J., Knuthsen, P., & Rasmussen, S. E. (2004). Urinary total flavonoid excretion but not 4-pyridoxic acid or potassium can be used as a biomarker for the intake of fruits and vegetables. *The Journal of Nutrition*, 134, 445–451.
- Landberg, R., Kamal-Eldin, A., Andersson, S.-O., Johansson, J.-E., Zhang, J. X., Hallmans, G., et al. (2009). Reproducibility of plasma alkylresorcinols during a 6-week rye intervention study in men with prostate cancer. *The Journal of Nutrition*, 139, 975–980.
- Lassale, C., Gunter, M. J., Romaguera, D., Peelen, L. M., Van der Schouw, Y. T., Beulens, J. W., et al. (2016). Diet quality scores and prediction of all-cause, cardiovascular and cancer mortality in a Pan-European cohort study. *PLoS One*, 11, e0159025.
- Linko-Parvinen, A. M., Landberg, R., Tikkanen, M. J., Adlercreutz, H., & Peñalvo, J. L. (2007). Alkylresorcinols from whole-grain wheat and rye are transported in human plasma lipoproteins. *Journal of Nutrition*, 137, 1137–1142.
- Lloyd, A. J., Beckmann, M., Favé, G., Mathers, J. C., & Draper, J. (2011). Proline betaine and its biotransformation products in fasting urine samples are potential biomarkers of habitual citrus fruit consumption. *British Journal of Nutrition*, 106, 812–824.
- Marklund, M., Magnusdóttir, O. K., Rosqvist, F., Cloetens, L., Landberg, R., Kolehmainen, M., et al. (2014). A dietary biomarker approach captures compliance and cardiometabolic effects of a healthy Nordic diet in individuals with metabolic syndrome. *The Journal of Nutrition*, 144, 1642–1649.
- McGrath, A. J., Hamill, L. L., Cardwell, C. R., Draffin, C. R., Neville, C. E., Appleton, K. M., et al. (2016). Combining vitamin C and carotenoid biomarkers better predicts fruit and vegetable intake than individual biomarkers in dietary intervention studies. *European Journal of Nutrition*, 55, 1377–1388.
- Monagas, M., Urpi-Sarda, M., Sánchez-Patán, F., Llorach, R., Garrido, I., Gómez-Cordovés, C., et al. (2010). Insights into the metabolism and microbial biotransformation of dietary flavan-3-ols and the bioactivity of their metabolites. *Food & Function*, 1, 233–253.
- Monošík, R., & Dragsted, L. O. (2016). A versatile UHPLC–MS/MS method for simultaneous quantification of various alcohol intake related compounds in human urine and blood. *Analytical Methods*, 8, 6865–6871.
- Neuhouser, M. L., Patterson, R. E., King, I. B., Horner, N. K., & Lampe, J. W. (2003). Selected nutritional biomarkers predict diet quality. *Public Health Nutrition*, 6, 703–709.
- Nielsen, S. E., Freese, R., Kleemola, P., & Mutanen, M. (2002). Flavonoids in human urine as biomarkers for intake of fruits and vegetables flavonoids in human urine as biomarkers for intake of fruits. *Cancer Epidemiology and Prevention Biomarkers*, 11, 459–466.
- O’Gorman, A., Morris, C., Ryan, M., O’Grada, C. M., Roche, H. M., Gibney, E. R., et al. (2014). Habitual dietary intake impacts on the lipidomic profile. *Journal of Chromatography B*, 966, 140–146.
- Pekkinen, J., Rosa-Sibakov, N., Micaud, V., Keski-Rahkonen, P., Lehtonen, M., Poutanen, K., et al. (2015). Amino acid-derived betaines dominate as urinary markers for rye bran intake in mice fed high-fat diet—A nontargeted metabolomics study. *Molecular Nutrition & Food Research*, 59, 1550–1562.
- Pujos-Guillot, E., Hubert, J., Martin, J.-F., Lyan, B., Quintana, M., Claude, S., et al. (2013). Mass spectrometry-based metabolomics for the discovery of biomarkers of fruit and vegetable intake: Citrus fruit as a case study. *Journal of Proteome Research*, 12, 1645–1659.
- Queipo-Ortuño, M. I., Boto-Ordóñez, M., Murri, M., Gomez-Zumaquero, J. M., Clemente-Postigo, M., Estruch, R., et al. (2012). Influence of red wine polyphenols and ethanol on the gut microbiota ecology and biochemical biomarkers. *The American Journal of Clinical Nutrition*, 95, 1323–1334.
- Rabassa, M., Cherubini, A., Zamora-Ros, R., Urpi-Sarda, M., Bandinelli, S., Ferrucci, L., et al. (2015). Low levels of a urinary biomarker of dietary polyphenol are associated with substantial cognitive decline over a 3-year period in older adults: The invecchiare in chianti study. *Journal of the American Geriatrics Society*, 63, 938–946.
- Rabassa, M., Zamora-Ros, R., Urpi-Sarda, M., Bandinelli, S., Ferrucci, L., Andres-Lacueva, C., et al. (2015). Association of habitual dietary resveratrol exposure with the development of frailty in older age: The invecchiare in chianti study. *The American Journal of Clinical Nutrition*, 102, 1534–1542.
- Rabassa, M., Zamora-Ros, R., Andres-Lacueva, C., Urpi-Sarda, M., Bandinelli, S., Ferrucci, L., et al. (2016). Association between both total baseline urinary and dietary polyphenols and substantial physical performance decline risk in older adults: A 9-year follow-up of the InCHIANTI study. *Journal of Nutrition, Health and Aging*, 20, 478–484.
- Rangel-Huerta, O. D., Aguilera, C. M., Perez-de-la-Cruz, A., Vallejo, F., Tomas-Barberan, G., Gil, A., et al. (2017). A serum metabolomics-driven approach predicts orange juice consumption and its impact on oxidative stress and inflammation in subjects from the BIONAOS study. *Molecular Nutrition & Food Research*, 61, 1600120.
- Rollo, M. E., Ash, S., Lyons-Wall, P., & Russell, A. (2011). Trial of a mobile phone method for recording dietary intake in adults with type 2 diabetes: Evaluation and implications for future applications. *Journal of Telemedicine and Telecare*, 17, 318–323.
- Rotches-Ribalta, M., Urpi-Sarda, M., Llorach, R., Boto-Ordóñez, M., Jauregui, O., Chiva-Blanch, G., et al. (2012). Gut and microbial resveratrol metabolite profiling after moderate long-term consumption of red wine versus dealcoholized red wine in humans by an optimized ultra-high-pressure liquid chromatography tandem mass spectrometry method. *Journal of Chromatography A*, 1265, 105–113.
- Rothwell, J. A., Fillâtre, Y., Martin, J. F., Lyan, B., Pujos-Guillot, E., Fezeu, L., et al. (2014). New biomarkers of coffee consumption identified by the non-targeted metabolomic profiling of cohort study subjects. *PLoS ONE*, 9, e93474.
- Scalbert, A., Brennan, L., Manach, C., Andres-Lacueva, C., Dragsted, L. O., Draper, J., et al. (2014). The food metabolome: A window over dietary exposure. *American Journal of Clinical Nutrition*, 99, 1286–1308.
- Schatzkin, A., Subar, A. F., Moore, S., Park, Y., Potischman, N., Thompson, F. E., et al. (2009). Observational epidemiologic studies of nutrition and cancer: The next generation (with better observation). *Cancer Epidemiology, Biomarkers & Prevention*, 18, 1026–1032.
- Selma, M. V., Beltrán, D., García-Villalba, R., Espín, J. C., & Tomás-Barberán, F. A. (2014). Description of urolithin production capacity from ellagic acid of two human intestinal Gordonibacter species. *Food & Function*, 5, 1779–1784.
- Shim, J. S., Oh, K., & Kim, H. C. (2014). Dietary assessment methods in epidemiologic studies. *Epidemiology and Health*, 36, e2014009.
- Slimani, N., Casagrande, C., Nicolas, G., Freisling, H., Huybrechts, I., Ocké, M. C., et al. (2011). The standardized computerized 24-h dietary recall method EPIC-Soft adapted for pan-European dietary monitoring. *European Journal of Clinical Nutrition*, 65, S5–S15.
- Stumbo, P. J. (2013). New technology in dietary assessment: A review of digital methods in improving food record accuracy. *The Proceedings of the Nutrition Society*, 72, 70–76.
- Sun, M., Fernstrom, J. D., Jia, W., Hackworth, S. A., Yao, N., Li, Y., et al. (2010). A wearable electronic system for objective dietary assessment. *Journal of the American Dietetic Association*, 110, 45–47.
- Tucker, K. L. (2007). Assessment of usual dietary intake in population studies of gene-diet interaction. *Nutrition, Metabolism, and Cardiovascular Diseases*, 17, 74–81.
- U.S. Department of Health and Human Services, & U.S. Department of Agriculture. (2015). *2015 – 2020 dietary guidelines for Americans* (8th ed., p. 2) December 2015. Available at: <http://health.gov/dietaryguidelines/2015/guidelines/>.
- Urpi-Sarda, M., Boto-Ordóñez, M., Queipo-Ortuño, M. I., Tulipani, S., Corella, D., Estruch, R., et al. (2015). Phenolic and microbial-targeted metabolomics to discovering and evaluating wine intake biomarkers in human urine and plasma. *Electrophoresis*, 36, 2259–2268.
- Varraso, R., Garcia-Aymerich, J., Monier, F., Le Moual, N., De Batlle, J., Miranda, G., et al. (2012). Assessment of dietary patterns in nutritional epidemiology: Principal component analysis compared with confirmatory factor analysis. *The American Journal of Clinical Nutrition*, 96, 1079–1092.
- Vázquez-Fresno, R., Llorach, R., Urpi-Sarda, M., Khymentis, O., Bulló, M., Corella, D., et al. (2015). An NMR metabolomics approach reveals a combined-biomarkers model in a wine interventional trial with validation in free-living individuals of the PREDIMED study. *Metabolomics*, 11, 797–806.
- Vermeulen, M., van Rooijen, H. J. M., & Vaes, W. H. J. (2003). Analysis of isothiocyanate mercapturic acids in urine: A biomarker for cruciferous vegetable intake. *Journal of Agricultural and Food Chemistry*, 51, 3554–3559.
- Zamora-Ros, R., Rabassa, M., Cherubini, A., Urpi-Sarda, M., Bandinelli, S., Ferrucci, L., et al. (2013). High concentrations of a urinary biomarker of polyphenol intake are associated with decreased mortality in older adults 1.2. *Journal of Nutrition*, 143, 1445–1450.
- Zamora-Ros, R., Urpi-Sarda, M., Lamuela-Raventós, R. M., Estruch, R., Martínez-González, M. A., Bulló, M., et al. (2009). Resveratrol metabolites in urine as a biomarker of wine intake in free-living subjects: The PREDIMED Study. *Free Radical Biology and Medicine*, 46, 1562–1566.